

# Upper ocean heat budget in the southeast Pacific stratus cloud region using eddy-resolving HYCOM

Yangxing Zheng (University of Colorado)

Toshi Shinoda (Naval Research Laboratory)

George Kiladis (NOAA-ESRL/PSD)


Jialin Lin (Ohio State University)

Joseph Metzger (Naval Research Laboratory)

Harley Hurlburt (Naval Research Laboratory)

Benjamin Geise (Texas A&M Univ.)

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>2009</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2009 to 00-00-2009</b>	
4. TITLE AND SUBTITLE <b>Upper ocean heat budget in the southeast Pacific stratus cloud region using eddy-resolving HYCOM</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Research Laboratory,Stennis Space Center,MS,39529</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>Layered Ocean Model Workshop (LOM 2009), Miami, FL</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>16</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			



*Stratus cloud decks ==> important role in regional and global climate*

*Most coupled GCMs have problems in producing realistic stratus clouds.*

*Upper ocean processes that control SST is crucial for simulating stratus clouds*

*Surface mooring measurements by WHOI since October 2000 (Colbo and Weller 2007)*

*New campaign: VAMOS Ocean-Cloud-Atmosphere-Land Study (VOCALS)*

*Intensive Observations: VOCALS Regional Experiment (Oct. 2008)*

## Upper ocean heat balance (upper 250m) at the buoy site (Colbo and Weller, 2007)

<i>Term</i>	<i>Estimate (W/m<sup>2</sup>)</i>	<i>Data</i>
<i>Surface heat flux</i>	<i>44 (± 5)</i>	<i>IMET buoy</i>
<i>Horizontal heat advection by Ekman transport</i>	<i>6 ( ± 4)</i>	<i>QuikSCAT winds Satellite SST</i>
<i>Horizontal heat advection by geostrophic transport</i>	<i>-20( ± 5)</i>	<i>IMET velocity, temperature Historical temperature</i>
<i>Ekman pumping</i>	<i>2.5( ± 5)</i>	<i>QuikSCAT winds IMET temperature</i>
<i>Eddy flux divergence</i>	<i>-30</i>	<i>Residual</i>
<i>Vertical diffusion</i>	<i>-3( ± 2)</i>	<i>IMET temperature</i>

## **Hypothesis (Colbo and Weller 2007)**

*“We postulate that the eddy flux divergence represents the effect of the cold coherent eddies formed near the coast..... the upwelled water does influence the offshore structure, but through the fluctuating mesoscale flow not the mean transport.”*

## **Issues**

*Upper ocean processes which balance the positive (warming) surface heat flux*

*Representativeness of the IMET site for the entire stratus cloud region*

*Role of eddies*

## **Models**

### ***Global HYCOM***

*Horizontal resolution: 1/12 deg. X 1/12 deg.*

*Period: Jan. 2003-Apr. 2007*

*No data assimilation*




### ***SODA***

*Horizontal resolution: 0.4 deg. (lon) X 0.25 deg.(lat)*


*Period: 1980-2005*

*Data assimilation*

*Relative importance of horizontal advection and eddy flux  
divergence for the upper ocean heat balance*



HYCOM: Net Surface Heat Flux






*Comparison of heat budget at 85W, 20S*


	<i>Colbo and Weller (2007)</i>	<i>HYCOM</i>	<i>SODA</i>
<i>Surface heat fllux</i>	44	18	
<i>Geostrophic</i>	-20	-45	-21
<i>Ekman</i>	6	-44	11
<i>Eddy flux div.</i>	-30	42	-19

*Ekman currents  $\rightarrow$  total - geostrophic*

## *Upper 50m*




## *Upper 50m*







SODA: Ekman Heat Flux




Sea Surface Eddy Kinetic Energy ((cm/s)<sup>2</sup>)








(a) SODA: Eddy Heat Divergence



(b) HYCOM: Eddy Heat Divergence



## Heat budget in the upper 50m (100W-80W, 30S-10S)

	<i>HYCOM</i>	<i>SODA</i>
<i>Surface heat flux</i>	<i>11</i>	
<i>Geostrophic</i>	<i>-5</i>	<i>-9</i>
<i>Ekman</i>	<i>3</i>	<i>6</i>
<i>Eddy flux div.</i>	<i>0.1</i>	<i>-0.1</i>



## Conclusions

*Geostrophic transport in the upper 50m causes net cooling in most of the stratus cloud region*

*Ekman transport provides net warming north of the IMET site and net cooling south of the IMET site*

*The eddy heat flux divergence term can be comparable to other terms at a particular location such as the IMET site, but it is negligible for the entire stratus region when area averaged since it is not spatially coherent in the open ocean.*

*Surface buoy observations in locations both north and south of the IMET site would be useful.*